•,								Sheet I U	<u>'                                    </u>
LIST OF REFERENCES CITED BY APPLICANT					ATTY DOCKET NO. APPLICATION NO 10873-008-999 09/492,764 APPLICANT				
(Use several sheets if necessary)				Jove et al.					
					FILING DATE GROUP 11 1642 5 1642 5 1642 5 1642				
U.S. PATENT DOCUMENTS									
*EXAMINER INITIAL		DOCUMENT NUMBER	DATE	и	АМЕ	CLASS	SUBCLASS;	FILING D IF APPROPI	ATE RIATE
KAC	CJ	5,935,993	8/10/99	Tang et al.		$\zeta$	5=	7	
KAC	CK	5,972,598	10/26/99	Chaudhary et al.			(5)		
AL	CL	6,426,366	7/30/02	Novogrodsky et al.		>	7	5	
FOREIGN PATENT DOCUMENTS									
		DOCUMENT NUMBER	DATE . COUNTRY CLASS			SUBCLASS	TRANSL		
		1	T			I	T	YES	NO
						<u> </u>		1	
	_								

		OTHER REFERENCES (Including Author, Title, Date, Pertinent Pages, Etc.)				
	СМ	Bright et al. IL-12 induced JAK-STAT pathways in T lymphocytes: Regulation by tyrphostin. J. of Allergy and				
		Clinical Immunology, 1997, 99(1 part 2):S287. Joint Meeting of the American Academy of Allergy, Asthma and				
KAC	ŀ	Immunology, the American Association of Immunologists and the Clinical Immunology Society; San Francisco, CA,				
100		USA; February 21-26, 1997				
	CN	Burger et al. IL-6 induced proliferation of a myeloma cell line is accompanied by activation of the JAK/STAT				
1		pathway and inhibited by tyrphostin AG490. Annals of Hematology, 1998, 77(suppl. 2):S2				
<del>     </del>	СО	Han et al. Preferential inhibition of glioblastoma cells with wild-type epidermal growth factor receptors by a novel				
1 /		tyrosine kinase inhibitor ethyl-2,5-dihydroxycinnamate. Oncol Res. 1997, 9(11-12):581-587				
	CP	Han et al. Tyrphostin AG 1478 preferentially inhibits human glioma cells expressing truncated rather than wild-type				
-		epidermal growth factor receptors. Cancer Res. 1996, 56(17):3859-3861				
	CQ	Heller et al. Treatment of cutaneous and subcutaneous tumors with electrochemotherapy using intralesional				
		bleomycin. DATABASE BIOSIS (online), Biosciences information service, Philadelphia, PA, US; July 1, 1998,				
		Databse accession No. PREV199800321817. Abstract & cancer, 83(1):148-157				
	CR	Lei et al. Enhancement of chemosensitivity and programmed cell death by tyrosine kinase inhibitors correlates with				
		EGFR expression in non-small cell lung cancer cells. Anticancer Res. 1999, 19(1A):221-228				
	CS	Liang et al. Chemosensitization of glioblastoma cells to bis-dichloroethyl-nitrosourea with tyrphostin AG17. Clin				
		Cancer Res. 1998, 4(3):773-781				
	CT	Nielsen et al. Constitutive activation of a slowly migrating isoform of Stat3 in mycosis fungoides: tyrphostin AG490				
		inhibits Stat3 activation and growth of mycosis fungoides tumor cell lines. Proc. Natl. Acad. Sci. U.S.A. 1997,				
		94(13):6764-6769				
	CU	Palumbo et al. The tryphostin AG17 induces apoptosis and inhibition of cdk2 activity in a lymphoma cell line that				
		overexpresses bcl-2. Cancer Res. 1997, 57(12):2434-2439				
	CV	Penar et al. Inhibition of epidermal growth factor receptor-associated tyrosine kinase blocks glioblastoma invasion of				
A		the brain. Neurosurgery. 1997, 40(1):141-151				

BY HAND DELIVERY
Shee A of 2

···		Sheery 01 2
1	CW	Tsai et al. Enhancement of chemosensitivity by tyrphostin AG825 in high-p185(neu) expressing non-small cell lung
KAC		cancer cells. Cancer Res. 1996, 56(5):1068-1074
	CX	Zushi et al. STAT3 mediates the survival signal in oncogenic ras-transfected intestinal epithelial cells. Int J Cancer.
		1998, 78(3):326-330
	CY	Zushi et al. Role of heparin-binding EGF-related peptides in proliferation and apoptosis of activated ras-stimulated
	]	intestinal epithelial cells. Int. J. Cancer 1997, 73(6):917-923
W		
		1

EXAMINER	Jarin	A.	Ganelle	DATE CONSIDERED	est 05/27/04
	/	<i>t</i>	7		